

AD-A043 036

MASSACHUSETTS INST OF TECH CAMBRIDGE DEPT OF CHEMISTRY F/G 7/3
PRESERVATION OF METAL-CARBON BONDS DURING DIRECT FLUORINATION. --ETC(U)
AUG 77 E K LIU, R J LAGOW
TR-2

N00014-75-C-0994
NL

UNCLASSIFIED

| OF |
AD
A043036



END
DATE
FILMED
9 -77
DDC

AD A 043036

OFFICE OF NAVAL RESEARCH

Contract N00014-75-C-0994
Task No. NR 053-601

Technical Report #2

PRESERVATION OF METAL-CARBON BONDS DURING
DIRECT FLUORINATION, A PRACTICAL SYNTHETIC METHOD.
THE SYNTHESIS OF TETRAKIS(TRIFLUOROMETHYL)GERMANIUM.

by

Edmund K. S. Liu and Richard J. Lagow

Department of Chemistry
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139



Contract checked cont. book

Reproduction in whole or in part is permitted for
any purpose of the United States Government.

SECTION for	
1115	White Section <input checked="" type="checkbox"/>
1115	Buff Section <input type="checkbox"/>
UNANNOUNCED <input type="checkbox"/>	
JUSTIFICATION	
DISTRIBUTION/AVAILABILITY CODES	
JIS. AVAIL. and/or SPECIAL	

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 2	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER 9
4. TITLE (and Subtitle) PRESERVATION OF METAL-CARBON BONDS DURING DIRECT FLUORINATION. A PRACTICAL SYNTHETIC METHOD. THE SYNTHESIS OF TETRAKIS(TRIFLUOROMETHYL)GERMANIUM.		5. TYPE OF REPORT & PERIOD COVERED Technical / Rept.
7. AUTHOR(s) Edmund K. S. Liu and Richard J. Lagow		6. PERFORMING ORG. REPORT NUMBER N00014-75-C-0994
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry University of Texas at Austin Austin, Texas 78712		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 053-601
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy Arlington, Virginia 22217		12. REPORT DATE 8/1/77
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 14 TR-2 12 1/2		13. NUMBER OF PAGES 5
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		15. SECURITY CLASS. (of this report) unclassified
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES To be published in the Journal of Chemical Society.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The metal-carbon bonds in tetramethylgermanium have been preserved during direct fluorination. Tetrakis(trifluoromethyl)germanium has been produced from the controlled reaction of elemental fluorine with tetramethyl-germanium.		

220 007

Contribution from the
Department of Chemistry
University of Texas at Austin
Austin, Texas 78712

Preservation of Metal-Carbon Bonds During Direct Fluorination, a Practical
Synthetic Method. The Synthesis of Tetrakis(trifluoromethyl)germanium.

by

E. Liu and R. J. Lagow *

Summary: The metal-carbon bonds in tetramethylgermanium have been preserved during direct fluorination. Tetrakis(trifluoromethyl)germanium has been produced from the controlled reaction of elemental fluorine with tetramethylgermanium.


Recently we have reported the first cases of preservation of mercury-carbon and silicon-carbon bonds during direct fluorination.¹ This synthetic method as applied to metal alkyls was regarded even in our laboratory as only a chemical curiosity. Recent results have shown that in some cases direct fluorination may be not only a practical synthetic method but one of choice for the preparation of perfluoroalkyl organo-metallics.

By controlling the fluorine concentration and temperature, we are able to obtain tetrakis(trifluoromethyl)germanium, $\text{Ge}(\text{CF}_3)_4$, in 63.5% yield from the reaction of tetramethylgermanium and fluorine. The experimental apparatus used has been discussed previously.² The physical and spectral properties of $\text{Ge}(\text{CF}_3)_4$ are identical with those of the same

compound first prepared in 1975 in our laboratory using plasma techniques.³

The best fluorination conditions are the following:

<u>Fluorine Flow</u>	<u>Helium Flow</u>	<u>Temperature</u>	<u>Hours</u>
1 cc/min	60 cc/min	-100 ⁰	48
"	"	-90	12
"	"	-80	14
"	"	-70	8
"	"	-60	14
"	"	-50	10
"	"	-40	12
"	"	-30	10
"	"	-20	14



The reaction is remarkably temperature dependent due to the extremely low reactivity of the partially fluorinated tetramethylgermanium compounds. If the reaction is run with a fluorine flow of 1 cc/minute to a helium flow of 60 cc/minute at -100⁰C for 150 hours, the yield of Ge(CF₃)₄ is only 1.1%. However if the reaction is run with a fluorine flow of 1 cc/minute to a helium flow of 60 cc/minute at -100⁰C for 2 days and the temperature increased 10⁰ per day to -60⁰, the yield of Ge(CF₃)₄ increases to 6%.

When the yield of Ge(CF₃)₄ is low, one obtains a number of partially fluorinated tetramethylgermanium compounds. Most of the compounds contain

Table 1^a

	¹ H ^b				¹⁹ F ^c							
	CH ₂ F	J _{HF}	CHF ₂	J _{HF}	CF ₃	J _{FF} ^d	CF ₂ H	J _{HF}	J _{FF}	CFH ₂	J _{HF}	J _{FF}
Ge(CF ₃) ₄					-27.0							
Ge(CF ₃) ₃ (CF ₂ H)			6.10	45.0	-27.2	3.0	49.0	45.5	3.1			
Ge(CF ₃) ₂ (CF ₂ H) ₂			6.23	45.5	-27.6	3.2	49.4	46.0	3.1			
Ge(CF ₃) ₂ (CF ₂ H)(CFH ₂)	4.98	46.5	6.24	45.7	-26.3	3.3	50.6	45.5	3.0	193.2	46.5	3.3
Ge(CF ₃)(CF ₂ H) ₃	6.25		6.25	45.5	-27.9	3.2	49.7	46.0	3.1			
Ge(CF ₃)(CF ₂ H) ₂ (CFH ₂)	4.89	46.0	6.15	45.5	-26.75	3.2	50.5	45.6	3.0	193.0	46.0	2.9
Ge(CF ₃)(CF ₂ H)(CFH ₂) ₂	4.90	46.0	6.25	45.6	-25.22	3.2	51.8	45.5	3.0	192.0	46.0	2.7
Ge(CF ₂ H) ₂ (CFH ₂) ₂	4.97	46.0	6.26	45.2			51.4	46.0	2.5	192.5	46.6	2.5

a All samples were run as neat liquids. Shifts are in parts per million. Coupling constants are in hertz.

b + downfield from external TMS

c + upfield from TFA

d The appropriate fluorine-fluorine coupling pattern present with n fluorine giving n+1 coupled pattern.

the appropriate integration are sufficient for identification of the compounds. For example, the compound $\text{Ge}(\text{CF}_3)_2(\text{CF}_2\text{H})_2$ gives the following NMR: H: CF_2H group; triplet, $J_{\text{HF}} = 45.5$; F: CF_3 group, pentet, $J_{\text{FF}} = 3.2$ resulting from coupling of two CF_2H groups, and CF_2H group, doublet, $J_{\text{HF}} = 46.0$, with each doublet being a septet, $J_{\text{FF}} = 3.1$ resulting from fluorine coupling of two CF_3 groups. The compounds were separated on 10% SE-30 on a chromsorb P columns.

ACKNOWLEDGEMENT

We are grateful for support from the Office of Naval Research for this work (N00014-75-C-0994).

References

1. E. Liu and R. J. Lagow, J. Amer. Chem. Soc., 98, 8270 (1976).
2. N. J. Maraschin, B. D. Catsikis, L. H. Davis, G. Jarvinen and R. J. Lagow, J. Amer. Chem. Soc., 97, 513 (1975).
3. R. J. Lagow, L. Gerchman, R. Jacob and J. Morrison, J. Amer. Chem. Soc., 97, 518 (1975).